


How do I apply SANTYL Ointment?

+ 4 simple steps to apply SANTYL Ointment



1 Cleanse

Gently cleanse the wound with sterile saline, or as directed by your health care professional.




2 Apply

Apply SANTYL Ointment directly to the wound at 2mm thickness. It should be about the **thickness of a nickel** lying flat.



3 Cover

Wounds with sufficient wound fluid will have enough moisture for product to be effective, but a dry wound may require additional moisture. Add moisture, as with a saline moistened gauze, compatible wound gel, and/or appropriate cleansers.



4 Change

Change dressings daily, or as instructed by your healthcare professional, to help remove dead tissue from the wound area to help advance the healing process.

Important Safety Information

SANTYL Ointment is used to remove damaged tissue from chronic skin ulcers and severely burned areas. Do not use if you have shown hypersensitivity to collagenase. Occasional slight redness may occur if SANTYL Ointment is placed outside the wound area. Precautions should be taken to avoid detergents or cleansing solutions that are acidic or basic in nature, as raising or lowering of the Ointment's pH can lower the effectiveness. Caregivers to debilitated patients should closely monitor the application site to ensure that the area is not becoming infected. The risk information provided here is not comprehensive. SANTYL[®] Collagenase Ointment - DIN 02063670

Advanced Wound Management

Smith & Nephew Inc.
2280 Argentia Road
Mississauga, ON
L5N 6H8 Canada
www.smith-nephew.com

Customer Care

T 1-800-463-7439
F 1-800-671-9140